

AD-A053 241


GEORGE WASHINGTON UNIV WASHINGTON D C INST FOR MANAG--ETC F/G 12/1
SENSITIVITY ANALYSIS IN NONLINEAR PROGRAMMING USING FACTORABLE --ETC(U)
MAR 78 A DE SILVA, G P MCCORMICK
N00014-75-C-0611
NL

UNCLASSIFIED

| OF |
AD
A053241



END
DATE
FILMED
6-78
DDC



12

AD A 053241

AD No.
DDC FILE COPY

THE
GEORGE
WASHINGTON
UNIVERSITY

STUDENTS FACULTY STUDY R
ESEARCH DEVELOPMENT FUT
URE CAREER CREATIVITY CO
MMUNITY LEADERSHIP TECH
NOLOGY FRONTIER DESIGN
ENGINEERING APP SCIENCE
GEORGE WASHINGTON UNIV

DDC
RECEIVED
APR 27 1978
B



INSTITUTE FOR MANAGEMENT
SCIENCE AND ENGINEERING
SCHOOL OF ENGINEERING
AND APPLIED SCIENCE

12

6

SENSITIVITY ANALYSIS IN NONLINEAR PROGRAMMING
USING FACTORABLE SYMBOLIC INPUT,

by

10

Anura/de Silva
Garth P./McCormick

12 21p.

14

Serial-T-365
8 March 1978

11 8 Mar 78

AD No. _____
DDC FILE COPY

The George Washington University
School of Engineering and Applied Science
Institute for Management Science and Engineering

Research Sponsored by

15

Office of Naval Research

Contract ~~NO000~~14-75-C-0611

Project NR 047 142

and

Federal Energy Administration

Purchase Order Number P-03-77-5938-0

DDC
RECEIVED
APR 27 1978
B

This document has been approved for public
sale and release; its distribution is unlimited.

406 743

LB

NONE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER T-365✓	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) SENSITIVITY ANALYSIS IN NONLINEAR PROGRAM- MING USING FACTORABLE SYMBOLIC INPUT		5. TYPE OF REPORT & PERIOD COVERED SCIENTIFIC
7. AUTHOR(s) ANURA de SILVA GARTH P. MCCORMICK		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS THE GEORGE WASHINGTON UNIVERSITY INSTITUTE FOR MANAGEMENT SCIENCE & ENGINEERING WASHINGTON, D. C. 20052		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0611 ^{NEW} FEA P.O. NO. P-03-77- 5938-0
11. CONTROLLING OFFICE NAME AND ADDRESS OFFICE OF NAVAL RESEARCH CODE 434 ARLINGTON, VIRGINIA 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE 8 March 1978
		13. NUMBER OF PAGES 19
		15. SECURITY CLASS. (of this report) NONE
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) SENSITIVITY ANALYSIS FACTORABLE PROGRAMMING NONLINEAR PROGRAMMING		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper develops the formulas and methodology to utilize the input to nonlinear programs in symbolic factorable form to perform first order sensitivity analysis on the solution vector of a general nonlinear program.		

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

NONE

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

THE GEORGE WASHINGTON UNIVERSITY
School of Engineering and Applied Science
Institute for Management Science and Engineering

Abstract
of
Serial T-365
8 March 1978

SENSITIVITY ANALYSIS IN NONLINEAR PROGRAMMING
USING FACTORABLE SYMBOLIC INPUT

by

Anura de Silva
Garth P. McCormick

This paper develops the formulas and methodology to utilize the input to nonlinear programs in symbolic factorable form to perform first order sensitivity analysis on the solution vector of a general nonlinear program.

Research Sponsored by
Office of Naval Research
and
Federal Energy Administration

ACCESSION for		
NTIS	White Section	<input checked="" type="checkbox"/>
DDC	Buff Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION _____		
BY _____		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
A		

THE GEORGE WASHINGTON UNIVERSITY
School of Engineering and Applied Science
Institute for Management Science and Engineering

SENSITIVITY ANALYSIS IN NONLINEAR PROGRAMMING
USING FACTORABLE SYMBOLIC INPUT

by

Anura de Silva
Garth P. McCormick

1. Introduction

This paper develops a methodology to utilize the input to nonlinear programs in symbolic factorable form to perform first order sensitivity analysis on the solution vector $x(y) \in E^n$ of a general nonlinear program, where $y \in E^k$ is the parametric vector. In Section 2 we discuss the key results of sensitivity analysis theory in nonlinear programming. In Section 3 we develop the formulae underlying the new methodology to utilize the symbolic factorable input to perform sensitivity analysis on the solution vector $x(y)$.

The above are implemented via the SUMT algorithm [4] for nonlinear programming which uses a penalty function technique. Section 4 briefly discusses the penalty function-based implementation of the sensitivity analysis methodology. An extrapolation result based on the penalty function parameter r is stated and incorporated in the computer code.

2. Relevant Results from NLP Sensitivity Theory

Consider the problem

$$\left. \begin{array}{l} \text{MIN} \quad F(x,y) \\ x \in E^n \\ \text{s.t.} \quad G(x,y) \geq 0 \\ \quad \quad H(x,y) = 0 \\ \text{where } G : E^n \times E^k \rightarrow E^m \\ \quad \quad H : E^n \times E^k \rightarrow E^p \end{array} \right\} \text{Problem } M(y)$$

By sensitivity information we mean the rates of change of the (locally) optimal objective function value and the corresponding optimizing vector with respect to changes in the parametric vector. If $x(y)$ is an n -vector and y a k -vector, the first order sensitivity information for Problem $M(y)$ is:

- (a) The rate of change of the solution value with respect to the parametric vector, $D_y F[x(y), y] \in E^k$.
- (b) The rate of change of the optimizing vector with respect to the parametric vector, $\frac{dx(y)}{dy}$, where

$$\frac{dx(y)}{dy} = \begin{bmatrix} \frac{\partial x_1(y)}{\partial y_1} & \dots & \frac{\partial x_1(y)}{\partial y_k} \\ \vdots & & \vdots \\ \frac{\partial x_n(y)}{\partial y_1} & \dots & \frac{\partial x_n(y)}{\partial y_k} \end{bmatrix} \quad (1)$$

The first order sensitivity results for the general nonlinear programming problem were first stated by Fiacco and McCormick in [3], Theorem 6, where they give conditions under which a smooth (once continuously differentiable, abbreviated OCD) trajectory of the optimizing Kuhn-Tucker triple $[x(y), u(y), w(y)]$ exist for y close to an initial value y_0 . The u and w are the Lagrange multiplier vectors associated with the inequality

constraints G and equality constraints H of Problem $M(y)$, in the Lagrangian,

$$L(x, y, u, w) = F(x, y) - \sum_{i=1}^m u_i G_i(x, y) + \sum_{j=1}^p w_j H_j(x, y) \quad (2)$$

where $u_i, G_i; i = 1, \dots, m$ and $w_j, H_j; j = 1, \dots, p$ are the components of the mappings G and H and the Lagrange multipliers u and w . Here, $u(y)$ and $w(y)$ are the values of these multipliers associated with the optimal value $x(y)$. They assume the following to hold; (a) twice continuous differentiability (abbreviated TCD) of $L(x, y, u, w)$ in (x, y) in a neighborhood of $(x(y_0), y_0)$, (b) second order sufficiency conditions (abbrev. SOSC) at $(x(y_0), y_0)$, (c) linear independence of binding constraints (abbrev. LIB) at $(x(y_0), y_0)$, (d) strict complementary slackness (abbrev. SCS). Here, (c) and (d) are regularity conditions they invoke to obtain the key result in a useful form. In a more recent paper Fiacco [2] obtains a similar result for somewhat more general parametric problem, such as $M(y)$.

The crux of the theory is the application of the implicit function theorem (see [6] for a general treatment of the implicit function theorem) to the first order Kuhn-Tucker system of $M(y)$, which is

$$\begin{aligned} \nabla_x L(x, y, u, w) &= 0 \\ u_i G_i(x, y) &= 0, \quad i = 1, \dots, m \\ H_j(x, y) &= 0, \quad j = 1, \dots, p. \end{aligned} \quad (3)$$

Consider a system of equations

$$D(x, y) = 0, \quad x \in E^n \quad \text{and} \quad y \in E^k \quad \text{Problem } P(y),$$

where $D : E^n \times E^k \rightarrow E^\ell$, ℓ being the number of scalar valued functions in D , which are assumed OCD. This ensures the existence of the Jacobians of $D(x, y)$ with respect to x and y , denoted $J_x D$ and $J_y D$, respectively. With the sensitivity analysis context in view, the implicit function theorem could be stated as follows:

Theorem 1. Suppose $D : E^n \times E^k \rightarrow E^l$; $(x(y_0), y_0)$ is a zero of $D(x, y)$; $D(x, y)$ is OCD near $(x(y_0), y_0)$; $J_x D$ is nonsingular at $(x(y_0), y_0)$.

Then, there exist open sets $S_x(x(y_0)) \subset E^n$ and $S_y(y_0) \subset E^k$ such that for any $y \in \bar{S}_y(y_0)$ there exists a unique solution for x in terms of y , i.e., $x = x(y) \in S_x(x(y_0))$, which is continuous in y . Moreover $x(y)$ is OCD at y_0 and

$$\nabla_y x(y_0) = - \left[J_x D_0 \right]^{-1} \left[J_y D_0 \right],$$

where the subscript 0 on D denotes evaluation at $(x(y_0), y_0)$.

If $P(y)$ is compared with (3) the role of Theorem 1 in the context of sensitivity analysis of the solution vector becomes evident. The assumed TCD of the problem functions gives the necessary OCD for the system in (3). The additional conditions of linear independence of the binding constraint gradients and SCS were assumed by Fiacco and McCormick [3] and Fiacco [2] to produce desirable results such as the uniqueness of the Lagrange multipliers $[u(y), w(y)]$ associated with the optimal $x(y)$ and the invariance of the set of inequality constraints binding at $x(y_0)$ in a neighborhood of y_0 . The following representation for the Kuhn-Tucker triple,

$$X \triangleq (x, u, w)$$

$$\text{and } X(y) \triangleq [x(y), u(y), w(y)]$$

will be adopted for notational convenience. For the same reason, the first order Kuhn-Tucker system of (3) will be denoted as follows.

$$R(x, u, w, y) = 0$$

or

$$\bar{R}(X, y) = 0.$$

(3b)

Differentiating (3b) partially with respect to X yields the Jacobian of (3b) or (3) with respect to X ,

$$J_X \bar{R}(X, y) = \begin{bmatrix} \nabla_{xx}^2 L & -\nabla_x G_1, \dots, -\nabla_x G_m & \nabla_x H_1, \dots, \nabla_x H_p \\ u_1 \nabla_x^T G_1 & G_1 & 0 \\ \vdots & \ddots & \vdots \\ u_m \nabla_x^T G_m & 0 & G_m \\ \nabla_x^T H_1 & & \\ \vdots & 0 & \\ \nabla_x^T H_p & & \end{bmatrix} \quad (4)$$

Differentiating (3b) partially with respect to y yields the Jacobian with respect to this parametric vector,

$$J_y \bar{R}(X, y) = \left[\nabla_{yx}^2 L^T, u_1 \nabla_y G_1, \dots, u_m \nabla_y G_m, \nabla_y H_1, \dots, \nabla_y H_p \right]^T. \quad (5)$$

At a Kuhn-Tucker point, (3b) could be written as

$$\bar{R}[X(y), y] = 0. \quad (3c)$$

Totally differentiating (3c) with respect to y , the following relationship is obtained.

$$J_X \bar{R}(X(y), y) \frac{dX(y)}{dy} + J_y \bar{R}(X(y), y) = 0. \quad (6)$$

The principal result immediately follows.

$$\frac{dX(y)}{dy} = - J_X \bar{R}(X(y), y)^{-1} \cdot J_y \bar{R}(X(y), y). \quad (7)$$

Equation (7) is almost the same as the result stated under Theorem 1 but is stated with respect to the entire Kuhn-Tucker triple, X . Note that (7) is stated for y in a neighborhood of y_0 , whereas the result of Theorem 1 was only at y_0 . This is possible due to the slightly stronger assumptions made by Fiacco and McCormick [3] and Fiacco [2].

The above result is the basis for the development of a methodology for sensitivity analysis in nonlinear programming. The computational technique used here is the penalty function method. The same result and technique were used by Armacost and Mylander [1] to compute (1), the principal difference in their methodology being that they approximate $J_y \bar{R}$ by central differencing techniques, and compute all other derivatives manually. The relevant formulae for the penalty function implementation using the logarithmic barrier-quadratic loss penalty function of the SUMT algorithm [4] are discussed in Section 3.

3. Sensitivity Analysis With Symbolic Input

The factorable input technique and computer code to compute gradient vectors and Hessian matrices of problem functions was developed by McCormick [5] to facilitate the use of nonlinear programming computer codes. The code itself has since been extended by de Silva and McCormick to accept input in a symbolic format. This symbolic code has been used to solve several general nonlinear programming problems and the ease with which a problem input can be supplied or modifications done is very encouraging. A sensitivity analysis option turned out to be a natural extension to the symbolic computer code.

A reduced version of the matrix in (1) which provides substantial computational economy, results from the formulae which are developed below; a brief discussion of the factorable input philosophy precedes this development.

In the factorable input method, the problem functions are built up recursively as sums, or sums of pairwise products, of simple transformations of previously defined functions. Initially, these simple transformations

are applied to the original problem variables x_i , $i = 1, \dots, n$, and the recursive application of the above procedure (shown mathematically in Equation (8)) generates new functions X^i , $i = n+1, \dots, N$.

McCormick [5] gives a detailed exposition illustrated by an example. He sets $X^i = x_i$, $i = 1, \dots, n$ and refers to the entire set of X^i , $i = 1, \dots, N$ as concomitant variable functions (abbrev. CVFs). The general equation for the creation of a new CVF is:

$$X^i(x) = \sum_{p=1}^{i-1} T_p^i[X^p(x)] + \sum_{p=1}^{i-1} \sum_{q=1}^p V_{q,p}^i[X^p(x)] \cdot U_{p,q}^i[X^q(x)] . \quad (8)$$

In the presence of a parametric vector y , the equation becomes

$$X^i(x,y) = \sum_{p=1}^{i-1} T_p^i[X^p(x,y),y] + \sum_{p=1}^{i-1} \sum_{q=1}^p V_{q,p}^i[X^p(x,y),y] \cdot U_{p,q}^i[X^q(x,y),y] . \quad (9)$$

The terms in the first summation of (8) or (9) are referred to as separable terms comprising the new CVF $X^i(\cdot)$ and the terms of the second summation of pairwise products are called quadratic terms. A point of note in (9) is that the arguments of the simple transformation

T_p^i , $V_{q,p}^i$ or $U_{p,q}^i$ contain y directly as well as indirectly via the

argument CVF. The direct presence of the parametric vector y is restricted to, at most, two components because all the simple transformations used in the methodology and code contain, at most, two constant terms. The parametric components are a subset of these constants.

An examination of (7) suggests that second partial derivatives of the problems functions with respect to x and the cross second partial derivatives with respect to x and y are required. This is because \bar{R} already contains first partial derivatives with respect to x . What is developed below will yield the matrix in (1) reduced by postmultiplying it

with a known column vector $\beta \in E^k$ or premultiplying it with a known row vector $\sigma \in E^n$. The postmultiplication produces an n-vector, each component being a weighted sum of the k partial derivatives of a particular $x_i(y)$, with respect to the components of the parametric vector.

The weights are the components of $\beta \in E^k$. The i th component of this resulting n-vector can be looked upon as the "total sensitivity" of the variable $x_i(y)$ ($i=1, \dots, n$) with respect to all the y_j , $j = 1, \dots, k$,

weighted according to β_j , $j = 1, \dots, k$. The r th component of the

n-vector is $\sum_{j=1}^k \frac{\partial x_r}{\partial y_j} \beta_j$. The premultiplication produces a k-component

vector, the j th component being the "total sensitivity" of a composite of all n variables $x_i(y)$, $i = 1, \dots, n$, weighted according to σ_i , $i = 1, \dots, n$, with respect to y_j ($j=1, \dots, k$). The ℓ th component of this k-vector

would be $\sum_{i=1}^n \frac{\partial x_i}{\partial y_\ell} \sigma_i$.

For clarity of exposition, (9) will be simplified below to a separable portion of one term or a quadratic term of one product.

Consider the separable case first. Suppose

$$X^i(x, y) = T[X^P(x, y), y] \quad (10)$$

Letting

$$\dot{T} \triangleq \dot{T}[X^P(x, y), y] \triangleq \frac{\partial T[X^P(x, y), y]}{\partial X^P}, \quad (11)$$

we have

$$\nabla_x X^i(x, y) = \nabla_x X^P(x, y) \cdot \frac{\partial T[X^P(x, y), y]}{\partial X^P} \quad (12a)$$

$$= \dot{T} \cdot \nabla_x X^P(x, y) \quad (12b)$$

In (12b) \dot{T} is readily available since all transformations are simple ones, and $\nabla_x X^P(x,y)$ is built up inductively from its argument CVF.

The expressions for $\nabla_{xx}^2 X^i(x,y)$ are already developed and coded in [5], and will not be repeated here. Note that

$$\nabla_{yx}^2 X^i(x,y) = \nabla_y \nabla_x X^i(x,y) \quad (13a)$$

$$= D_y D_x T[X^P(x,y),y] \quad (13b)$$

In (13b) D denotes total differentiation, although in the previous step the partial differentiation ∇ was used. This is due to the fact that the partial differentiation in (13a) would capture the total variation due to x and y of the expression in (12a), but not so in (13b). Continuing the differentiation with respect to y of (12b), as in (13b), we get

$$\begin{aligned} \nabla_{yx}^2 X^i(x,y) &= \nabla_{yx}^2 X^P(x,y) \cdot \dot{T} + \nabla_x X^P(x,y) \left\{ \dot{T} [X^P(x,y),y] \nabla_y^T X^P(x,y) \right. \\ &\quad \left. + \nabla_y^T \dot{T} \right\}. \end{aligned} \quad (14)$$

$$\text{In the above, } \ddot{T} \triangleq \dot{T} [X^P(x,y),y] \triangleq \frac{\partial^2 T[X^P(x,y),y]}{\partial (X^P)^2}. \quad (15)$$

In (14), \dot{T} and \ddot{T} are readily available scalar quantities. The quantity $\nabla_y^T \dot{T} [X^P(x,y),y]$ is the true partial derivative unlike in (13b). This is easy to compute because y enters the \dot{T} argument directly, at most, in two nonzero components. The expression for $\nabla_x X^P(x,y)$ has been derived already in (12a). The only remaining terms are $\nabla_{yx}^2 X^P(x,y)$ which is built up inductively, and $\nabla_y^T X^P(x,y)$ which is discussed below.

Now

$$\begin{aligned}\nabla_y X^i(x,y) &= D_y T[X^P(x,y),y] \\ &= \nabla_y T[X^P(x,y),y] + \dot{T} \cdot \nabla_y X^P(x,y) .\end{aligned}\tag{16}$$

Note $\nabla_y T(X^P(x,y),y)$ is the true partial derivative and will have at most two nonzero components in E^k which are easily computable. Thus (16) permits the computation of the $\nabla_y^T X^P(x,y)$ term of (14) by induction. Since the formulae of (12b), (14), (16) are initially applied to the original problem variables $X_i = x^i$, $i = 1, \dots, n$, we have, for $p \leq n$:

$$\nabla_x X^i(x,y) = e_p \cdot \dot{T} \tag{17}$$

where e_j is a unit vector $\in E^n$ with unity in the j th position.

$$\begin{aligned}\text{Also, } \nabla_{yx}^2 X^i(x,y) &= \nabla_{yx}^2 X^P + e_p [\dot{T} \nabla^T X^P + \nabla_y^T \dot{T}] \\ &= e_p \nabla_y^T \dot{T} .\end{aligned}\tag{18}$$

Finally,

$$\begin{aligned}\nabla_y X^i(x,y) &= \nabla_y T + \nabla_y X^P \dot{T} \\ &= \nabla_y T .\end{aligned}\tag{19}$$

On page 6, we spoke of dimensionally reducing the matrix of (1) in two separate ways. The case of premultiplying by $\alpha \in E^n$ is considered below.

Thus, from (17) and (18) letting α_p denote the p th component of α , we get

$$\alpha^T \nabla_x X^i(x,y) = \alpha_p \cdot \dot{T} . \tag{20}$$

$$\alpha^T \nabla_{yx}^2 X^i(x,y) = \alpha_p \nabla_y^T \dot{T} . \tag{21}$$

The summation of (21) will have, at most, two nonzero components due to the nature of the simple transformations used. Next, consider the case of $p > n$. Here,

$$\alpha^T \nabla_x X^i(x,y) = \alpha^T \nabla_x X^p(x,y) \cdot \dot{T} \quad (22)$$

Next,

$$\begin{aligned} \alpha^T \nabla_{yx}^2 X^i(x,y) &= \alpha^T \nabla_{yx}^2 X^p(x,y) \cdot \dot{T} \\ &+ \alpha^T \nabla_x X^p(x,y) \left\{ \dot{T} \nabla_y^T X^p(x,y) + \nabla_y^T \dot{T} \right\} \end{aligned} \quad (23)$$

Equation (22) works purely inductively and does not require the deliberate introduction of α , as this has been done in (20). It is a scalar quantity for each i . The first term on the right-hand side of (23) is also an inductive term. As far as the second term is concerned, (the scalar multiplier $\alpha^T \nabla_x X^p(x,y)$, the only presence of α there,) it is built up inductively. The expression within the curly bracket is a k -vector, the first term being an inductive one as demonstrated in (16) and the second possessing, at most, two nonzero components.

In summary, we have a scalar quantity for the reduced version of the gradient vector of each CVF i , and (20) (or (17), if $p \leq n$) has to be repeated for every p corresponding to the separable terms of X^i , and added together. The reduced second cross partial of each CVF i (as shown in (21) or (18) depending on whether $p \geq n$) is a k -vector for each component p , which must be added together as before.

The quadratic components of (9) will be addressed next. Considering the creation of an X^i , $i = n+1, \dots, N$, using a single product, we have

$$X^i(x,y) = V[X^p(x,y), y] * U[X^q(x,y), y] \quad (24)$$

then,

$$\begin{aligned} \nabla_x X^i(x,y) &= D_x V[X^p(x,y), y] * U[X^q(x,y), y] \\ &+ V[X^p(x,y), y] * D_x U[X^q(x,y), y] \end{aligned} \quad (25)$$

$$= \dot{V}U \cdot \nabla_x X^p(x,y) + V\dot{U} \cdot \nabla_x X^q(x,y) \quad (26)$$

Now \dot{V} , \dot{U} are defined analogously to \dot{T} (see (11)). Since V , \dot{V} , U , \dot{U} are scalar quantities, the premultiplication by $\alpha \in E^n$ could be done as in (20) and (22) depending on whether p and q are \leq or $> n$. The separate terms in (25) must be added together. The result is

$$\alpha^T \nabla_x^i X^i(x, y) = \dot{V} U \alpha^T \nabla_x X^p(x, y) + V \dot{U} \alpha^T \nabla_x X^q(x, y). \quad (26)$$

When p (or q) is $> n$, $\alpha^T \nabla_x X^p$ (or $\alpha^T \nabla_x X^q$) enters purely inductively. Otherwise, the term simplifies to $\dot{V} U \alpha_p$ (or $V \dot{U} \alpha_q$). The second cross partial derivative is obtained by differentiating (26) with respect to y . The arguments of X^p and X^q are omitted for convenience of representation. Finally,

$$\begin{aligned} \nabla_{yx}^2 X^i(x, y) = U \left\{ \dot{V} \cdot \nabla_{yx}^2 X^p + \nabla_x X^p [\dot{V} \cdot \nabla_y^T X^p + \nabla_y^T \dot{V}] \right\} \\ + \dot{V} \cdot \nabla_x X^p \cdot D_y U \\ + V \left\{ \dot{U} \cdot \nabla_{yx}^2 X^q + \nabla_x X^q [\dot{U} \cdot \nabla_y^T X^q + \nabla_y^T \dot{U}] \right\} \\ + \dot{U} \cdot \nabla_x X^q \cdot D_y V. \end{aligned} \quad (27)$$

Premultiplying (27) by $\alpha \in E^n$, and rearranging,

$$\begin{aligned} \alpha^T \nabla_{yx}^2 X^i(x, y) = U \left\{ \alpha^T \nabla_{yx}^2 X^p \cdot \dot{V} + \alpha^T \nabla_x X^p [\dot{V} \cdot \nabla_y^T X^p + \nabla_y^T \dot{V}] \right\} \\ + V \left\{ \alpha^T \nabla_{yx}^2 X^q \cdot \dot{U} + \alpha^T \nabla_x X^q [\dot{U} \cdot \nabla_y^T X^q + \nabla_y^T \dot{U}] \right\} \\ + \alpha^T \nabla_x X^p \cdot D_y U \cdot \dot{V} + \alpha^T \nabla_x X^q \cdot \nabla_y D \cdot \dot{U}. \end{aligned} \quad (28)$$

The evaluation of the portions within the curly brackets in (28) is identical to the evaluation of the cross second partials in the separable analysis, i.e., the right-hand side of (23). The last two terms of (26) contain two quantities $\alpha^T \nabla_x X^p$ and $\alpha^T \nabla_x X^q$ which are developed by induction as demonstrated by (12) and (20). The quantities $D_y U$ and $D_y V$

are analogous to D_y^T whose computation is shown in (16). The only remaining expression to be developed is that for $\nabla_y X^i$,

$$\begin{aligned}\nabla_y X^i(x,y) &= D_y \{V[X^P(x,y),y] U[X^Q(x,y),y]\} \\ &= U\{\dot{V} \cdot \nabla_y X^P + \nabla_y V\} + V\{\dot{U} \cdot \nabla_y X^Q + \nabla_y U\}.\end{aligned}\quad (29)$$

Examination of (23) and (28) shows that it is not necessary to premultiply the expressions of (16) and (29) by α^T , since $\nabla_y X$ never gets directly premultiplied by it.

The above Equations (20) through (29) have developed the necessary components to perform the computations necessary to produce a row k-vector $\alpha^T [dx(y)/dy]$. The next section describes how these are implemented via a penalty function method.

4. Computation of Sensitivity Results

The SUMT computer code [4] is used in conjunction with the factorable symbolic sensitivity analysis code to yield $[\alpha^T dx(y)/dy]$. The code uses an interior-exterior penalty function with a single parameter r , and generates a sequence of subproblems which will be denoted as $M(y,r)$, which tend to $M(y)$ as $r \rightarrow 0$. In straight nonlinear optimization SUMT uses an extrapolation based on r to estimate the relevant values at $r = 0$. A similar extrapolation technique is used here to provide first order estimates of $[\alpha^T dx(y)/dy]$.

A logarithmic barrier quadratic loss penalty function is considered below. For $M(y)$, this penalty function is

$$P[x,y,r] = F(x,y) - r \sum_{i=1}^m \ln G_i(x,y) + \sum_{j=1}^p \frac{H_j^2(x,y)}{r}.\quad (30)$$

The problem solved at the ℓ th stage of the sequential procedure, for a value $r = r^\ell$ is

$$\begin{array}{ll}\text{MIN}_{x \in R^0} & P[x,y,r^\ell] \quad \text{Problem } M(r,y)\end{array}$$

where

$$R^0 \equiv \{x | G_i(x,y) > 0, i=1, \dots, m\}.$$

The optimality condition is that the gradient with respect to x vanishes, i.e.,

$$\nabla_x P[x, y, r^k] = 0. \quad (31)$$

At local optima, for a general value of r we can write

$$\nabla_x P[x(y, r), y, r] \equiv 0. \quad (32)$$

Applying this to (30),

$$\begin{aligned} \nabla_x F[x(y, r), y] - \sum_{i=1}^m \frac{r}{G_i[x(y, r), y]} \cdot \nabla_x G_i[x(y, r), y] \\ + \sum_{j=1}^p \frac{2 H_j[x(y, r), y]}{r} \nabla_x H_j[x(y, r), y] \equiv 0. \end{aligned} \quad (33)$$

Differentiating (33) partially with respect to y , we get

$$\begin{aligned} \nabla_{yx}^2 P[x(y, r), y, r] \\ = \nabla_{yx}^2 F[x(y, r), y] - \sum_{i=1}^m \nabla_{yx}^2 G_i[x(y, r), y] \cdot \frac{r}{G_i[x(y, r), y]} \\ + \sum_{i=1}^m \frac{r}{G_i[x(y, r), y]^2} \cdot \nabla_x G_i[x(y, r), y] \cdot \nabla_y^T G_i[x(y, r), y] \\ + \sum_{j=1}^p \nabla_{yx}^2 H_j[x(y, r), y] \cdot \frac{2 H_j[x(y, r), y]}{r} \\ + \sum_{j=1}^p \left(\frac{2}{r} \right) \nabla_x H_j[x(y, r), y] \cdot \nabla_y^T H_j[x(y, r), y]. \end{aligned} \quad (34)$$

Differentiating (33) totally with respect to y ,

$$\nabla_{xx}^2 P[x(y, r), y, r] \frac{dx(y, r)}{dy} + \nabla_{yx}^2 P[x(y, r), y, r] \equiv 0. \quad (35)$$

The argument variables within the square brackets will be omitted when referring to the terms in the above.

$\nabla_{xx}^2 P$ has been developed elsewhere by McCormick [5]. Those for $\nabla_{xy}^2 P$ are developed using the Equation (34). Thus,

$$\frac{d(x(y,r))}{dy} = - \nabla_{xx}^2 P[x(y,r), y, r]^{-1} \cdot \nabla_{yx}^2 P[x(y,r), y, r] . \quad (36)$$

The similarity between (7) and (36) is to be noted. Suppose (36) is premultiplied by some $\sigma \in E^n$. Then, we get

$$\sigma^T \frac{dx(y,r)}{dy} = - \alpha^T \cdot \nabla_{yx}^2 P[x(y,r), y, r] , \quad (37)$$

where

$$\alpha^T = \sigma^T \cdot \nabla_{xx}^2 P[x(y,r), y, r]^{-1} . \quad (38)$$

If σ is a known vector, α is determined because the Hessian of the penalty function with respect to x is known by [5]. Then (37) is determined by premultiplying (34) by this α , and using (20), (21), (22), (23), (27) and (29) to determine the individual components of the resulting equation. This can be done because every problem function F , G_i ; $i = 1, \dots, m$, H_j ; $j = 1, \dots, p$, can be identified with a CVF X^i , $i = 1, \dots, n$, $n+1, \dots, N$.

The final aspect remaining to be discussed in the context of the sensitivity methodology is the improvement of the estimate of $\sigma^T \frac{dx(y,r)}{dy}$ by extrapolating on r , at the ℓ th sequential unconstrained problem, to $r = 0$. A Taylor series approximation applied at the ℓ th and $(\ell-1)$ th subproblems yields

$$\frac{dx(y, r^l)}{dy} = \frac{dx(y, 0)}{dy} + r^l \frac{d}{dr} \left[\frac{dx(y, 0)}{dy} \right]. \quad (39)$$

$$\frac{dx(y, r^{l-1})}{dy} = \frac{dx(y, 0)}{dy} + r^{l-1} \frac{d}{dr} \left[\frac{dx(y, 0)}{dy} \right]. \quad (40)$$

Eliminating the last term in (39) and (40)

$$\frac{d(x(y, 0))}{dy} (r^{l-1} - r^l) = r^{l-1} \cdot \frac{dx(y, r^l)}{dy} - r^l \cdot \frac{dx(y, r^{l-1})}{dy} \quad (41)$$

from which, (letting $r^{l-1} = c \cdot r^l$) we get

$$\frac{dx(y, 0)}{dy} = \frac{1}{c-1} \cdot \left[c \frac{dx(y_1 r^l)}{dy} - \frac{dx(y_1 r^{l-1})}{dy} \right]. \quad (42)$$

Hence by performing the computation for successive subproblems when r becomes sufficiently small, (42) can be used to estimate the value for the original problem, i.e., $r = 0$.

REFERENCES

- [1] ARMACOST, R. L. and W. C. MYLANDER (1973). A guide to a SUMT-Version 4 computer subroutine for implementing sensitivity analysis in nonlinear programming. Technical Paper Serial T-237. The Institute for Management Science and Engineering, The George Washington University.
- [2] FIACCO, A. V. (1976). Sensitivity analysis for nonlinear programming using penalty methods. Mathematical Programming 10 (3) 287-311.
- [3] FIACCO, A. V. and G. P. MCCORMICK (1968). Nonlinear Programming: Sequential Unconstrained Minimization Techniques. Wiley, New York.
- [4] HOLMES, R. L., W. C. MYLANDER, and G. P. MCCORMICK (1971). A guide to SUMT-Version 4: The computer program implementing the sequential unconstrained minimization technique for nonlinear programming. RAC-P-63, Research Analysis Corporation, McLean, Virginia.
- [5] MCCORMICK, G. P. (1974). A mini-manual for use of the SUMT computer program and the factorable programming language. Technical Report SOL 74-15. Systems Optimization Laboratory, Stanford University.
- [6] ORTEGA, J. M. and W. C. RHEINBOLDT (1970). Iterative Solutions of Nonlinear Equations in Several Variables. Academic Press.

THE GEORGE WASHINGTON UNIVERSITY

Program in Logistics

Distribution List for Technical Papers

<p>The George Washington University Office of Sponsored Research Library Vice President H. F. Bright Dean Harold Liebowitz Mr. J. Frank Doubleday</p> <p>ONR Chief of Naval Research (Codes 200, 430D, 1021P) Resident Representative</p> <p>OPNAV OP-40 DCNO, Logistics Navy Dept Library OP-911 OP-964</p> <p>Naval Aviation Integrated Log Support</p> <p>NAVCOSSACT</p> <p>Naval Cmd Sys Sup Activity Tech Library</p> <p>Naval Electronics Lab Library</p> <p>Naval Facilities Eng Cmd Tech Library</p> <p>Naval Ordnance Station Louisville, Ky. Indian Head, Md.</p> <p>Naval Ordnance Sys Cmd Library</p> <p>Naval Research Branch Office Boston Chicago New York Pasadena San Francisco</p> <p>Naval Research Lab Tech Info Div Library, Code 2029 (ONRL)</p> <p>Naval Ship Engng Center Philadelphia, Pa. Hyattsville, Md.</p> <p>Naval Ship Res & Dev Center</p> <p>Naval Sea Systems Command Tech Library Code 073</p> <p>Naval Supply Systems Command Library Capt W. T. Nash</p> <p>Naval War College Library Newport</p> <p>BUPERS Tech Library</p> <p>FMSO</p> <p>Integrated Sea Lift Study</p> <p>USN Ammo Depot Earle</p> <p>USN Postgrad School Monterey Library Dr. Jack R. Borsting Prof C. R. Jones</p> <p>US Marine Corps Commandant Deputy Chief of Staff, R&D</p> <p>Marine Corps School Quantico Landing Force Dev Ctr Logistics Officer</p> <p>Armed Forces Industrial College</p> <p>Armed Forces Staff College</p> <p>Army War College Library Carlisle Barracks</p> <p>Army Cmd & Gen Staff College</p> <p>US Army HQ LTC George L. Slyman Army Trans Mat Command</p>	<p>Army Logistics Mgmt Center Fort Lee</p> <p>Commanding Officer, USALDSRA New Cumberland Army Depot</p> <p>US Army Inventory Res Ofc Philadelphia</p> <p>HQ, US Air Force AFADS-3</p> <p>Griffiss Air Force Base Reliability Analysis Center</p> <p>Maxwell Air Force Base Library</p> <p>Wright-Patterson Air Force Base HQ, AF Log Command Research Sch Log</p> <p>Defense Documentation Center</p> <p>National Academy of Science Maritime Transportation Res Board Library</p> <p>National Bureau of Standards Dr E. W. Cannon Dr Joan Rosenblatt</p> <p>National Science Foundation</p> <p>National Security Agency</p> <p>WSEG</p> <p>British Navy Staff</p> <p>Logistics, OR Analysis Establishment National Defense Hdqtrs, Ottawa</p> <p>American Power Jet Co George Chernowitz</p> <p>ARCON Corp</p> <p>General Dynamics, Pomona</p> <p>General Research Corp Dr Hugh Cole Library</p> <p>Planning Research Corp Los Angeles</p> <p>Rand Corporation Library</p> <p>Carnegie-Mellon University Dean H. A. Simon Prof G. Thompson</p> <p>Case Western Reserve University Prof B. V. Dean Prof John R. Isbell Prof M. Mesarovic Prof S. Zacks</p> <p>Cornell University Prof R. E. Bechhofer Prof R. W. Conway Prof J. Kiefer Prof Andrew Schultz, Jr.</p> <p>Cowles Foundation for Research Library Prof Herbert Scarf Prof Martin Shubik</p> <p>Florida State University Prof R. A. Bradley</p> <p>Harvard University Prof K. J. Arrow Prof W. G. Cochran Prof Arthur Schleifer, Jr.</p> <p>New York University Prof O. Morgenstern</p> <p>Princeton University Prof A. W. Tucker Prof J. W. Tukey Prof Geoffrey S. Watson</p>
---	--

Purdue University
Prof S. S. Gupta
Prof H. Rubin
Prof Andrew Whinston

Stanford
Prof T. W. Anderson
Prof G. B. Dantzig
Prof F. S. Hillier
Prof D. L. Iglehart
Prof Samuel Karlin
Prof G. J. Lieberman
Prof Herbert Solomon
Prof A. F. Veinott, Jr.

University of California, Berkeley
Prof R. E. Barlow
Prof D. Gale
Prof Rosedith Sitgreaves
Prof L. M. Tichvinsky

University of California, Los Angeles
Prof J. R. Jackson
Prof Jacob Marschak
Prof R. R. O'Neill
Numerical Analysis Res Librarian

University of North Carolina
Prof W. L. Smith
Prof M. R. Leadbetter

University of Pennsylvania
Prof Russell Ackoff
Prof Thomas L. Saaty

University of Texas
Prof A. Charnes

Yale University
Prof F. J. Anscombe
Prof I. R. Savage
Prof M. J. Sobel
Dept of Admin Sciences

Prof Z. W. Birnbaum
University of Washington

Prof B. H. Bissinger
The Pennsylvania State University

Prof Seth Bolder
University of Michigan

Prof G. E. P. Box
University of Wisconsin

Dr. Jerome Bracken
Institute for Defense Analyses

Prof H. Chernoff
MIT

Prof Arthur Cohen
Rutgers - The State University

Mr Wallace M. Cohen
US General Accounting Office

Prof C. Derman
Columbia University

Prof Paul S. Dwyer
Mackinaw City, Michigan

Prof Saul I. Gass
University of Maryland

Dr Donald P. Gaver
Carmel, California

Dr Murray A. Geisler
Logistics Mgmt Institute

Prof J. F. Hannan
Michigan State University

Prof H. O. Hartley
Texas A & M Foundation

Mr Gerald F. Hein
NASA, Lewis Research Center

Prof W. M. Hirsch
Courant Institute

Dr Alan J. Hoffman
IBM, Yorktown Heights

Dr Rudolf Husser
University of Bern, Switzerland

Prof J. H. K. Kao
Polytech Institute of New York

Prof W. Kruskal
University of Chicago

Prof C. E. Lemke
Rensselaer Polytech Institute

Prof Loynes
University of Sheffield, England

Prof Steven Nahmias
University of Pittsburgh

Prof D. B. Owen
Southern Methodist University

Prof E. Parzen
State University New York, Buffalo

Prof H. O. Posten
University of Connecticut

Prof R. Remage, Jr.
University of Delaware

Dr Fred Rigby
Texas Tech College

Mr David Rosenblatt
Washington, D. C.

Prof M. Rosenblatt
University of California, San Diego

Prof Alan J. Rowe
University of Southern California

Prof A. H. Rubenstein
Northwestern University

Dr M. E. Salvesson
West Los Angeles

Prof Edward A. Silver
University of Waterloo, Canada

Prof R. M. Thrall
Rice University

Dr S. Vajda
University of Sussex, England

Prof T. M. Whitin
Wesleyan University

Prof Jacob Wolfowitz
University of Illinois

Mr Marshall K. Wood
National Planning Association

Prof Max A. Woodbury
Duke University

THE GEORGE WASHINGTON UNIVERSITY

BENEATH THIS PLAQUE
IS BURIED
A VAULT FOR THE FUTURE
IN THE YEAR 2036

THE STORY OF ENGINEERING IN THIS YEAR OF THE PLACING OF THE VAULT AND
ENGINEERING HOPES FOR THE TOMORROWS AS WRITTEN IN THE RECORDS OF THE
FOLLOWING GOVERNMENTAL AND PROFESSIONAL ENGINEERING ORGANIZATIONS AND
THOSE OF THIS GEORGE WASHINGTON UNIVERSITY.

BOARD OF COMMISSIONERS DISTRICT OF COLUMBIA
UNITED STATES ATOMIC ENERGY COMMISSION
DEPARTMENT OF THE ARMY UNITED STATES OF AMERICA
DEPARTMENT OF THE NAVY UNITED STATES OF AMERICA
DEPARTMENT OF THE AIR FORCE UNITED STATES OF AMERICA
NATIONAL ADVISORY COMMITTEE FOR AERONAUTICS
NATIONAL BUREAU OF STANDARDS U.S. DEPARTMENT OF COMMERCE
AMERICAN SOCIETY OF CIVIL ENGINEERS
AMERICAN INSTITUTE OF ELECTRICAL ENGINEERS
THE AMERICAN SOCIETY OF MECHANICAL ENGINEERS
THE SOCIETY OF AMERICAN MILITARY ENGINEERS
AMERICAN INSTITUTE OF MINING & METALLURGICAL ENGINEERS
DISTRICT OF COLUMBIA SOCIETY OF PROFESSIONAL ENGINEERS, INC.
THE INSTITUTE OF RADIO ENGINEERS, INC.
THE CHEMICAL ENGINEERS CLUB OF WASHINGTON
WASHINGTON SOCIETY OF ENGINEERS
FAULKNER KINGSBURY & STENHOUSE - ARCHITECTS
CHARLES H. TOMPKINS COMPANY - BUILDERS
SOCIETY OF WOMEN ENGINEERS
NATIONAL ACADEMY OF SCIENCES NATIONAL RESEARCH COUNCIL

THE PURPOSE OF THIS VAULT IS INSPIRED BY AND IS DEDICATED TO
CHARLES HOOK TOMPKINS, DOCTOR OF ENGINEERING
BECAUSE OF HIS ENGINEERING CONTRIBUTIONS TO THIS UNIVERSITY, TO HIS
COMMUNITY, TO HIS NATION, AND TO OTHER NATIONS.

BY THE GEORGE WASHINGTON UNIVERSITY.

ROBERT V. FLEMING
CHAIRMAN OF THE BOARD OF TRUSTEES

CLOYD H. MARVIN
PRESIDENT

JUNE THE TWENTIETH
1956

To cope with the expanding technology, our society must be assured of a continuing supply of rigorously trained and educated engineers. The School of Engineering and Applied Science is completely committed to this objective.